

Integer powers of complex anti-tridiagonal matrices and some complex factorizations

Durmuş Bozkurt* & H. Kübra Duru†

Department of Mathematics, Science Faculty of Selçuk University

June 13, 2014

Abstract

In this paper, we obtain a general expression for the entries of the r th power of a certain $n \times n$ complex anti-tridiagonal matrix where if n is odd, $r \in \mathbb{Z}$ or if n is even, $r \in \mathbb{N}$. In addition, we get the complex factorizations of Fibonacci polynomials, Fibonacci and Pell numbers.

1 Introduction

Arbitrary integer powers of a square matrix is used to solve some difference equations, differential and delay differential equations and boundary value problems.

Recently, the calculations eigenvalues and integer powers of anti-tridiagonal matrices have been well studied in the literature. For instance, Rimas [1-3] obtained the integer powers of anti-tridiagonal matrices of odd and even order. Öteleş and Akbulak [8-10] generalized Rimas's the some results and obtained complex factorizations. Gutiérrez [11] calculated the powers of complex persymmetric or skew-persymmetric anti-tridiagonal matrices with costant anti-diagonals. For details on the eigenvalues and the powers of tridiagonal and anti-tridiagonal matrices, see [4-7, 12]. We obtained integer powers of the tridiagonal matrix

$$\tilde{A}_n = \begin{cases} \tilde{a}_{11} = \tilde{a}_{nn} = a \\ \tilde{a}_{12} = \tilde{a}_{n,n-1} = 2b \\ \tilde{a}_{21} = \tilde{a}_{n-1,n} = b \\ tridiag(-b, a, -b), \text{ other} \end{cases} \quad (1)$$

as

*Corresponding author, dbozkurt@selcuk.edu.tr

†hkdu@selcuk.edu.tr

$$\begin{aligned}\tilde{a}_{ij}(r) &= \frac{1}{2n-2} (\lambda_2^r T_{i-1}(m_2) T_{j-1}(m_2) + \lambda_3^r T_{i-1}(m_3) T_{j-1}(m_3)) \\ &\quad + 2 \sum_{\substack{k=1 \\ k \neq 2,3}}^n \lambda_k^r T_{i-1}(m_k) T_{j-1}(m_k); \quad i = 1, \dots, n; \quad j = 1, n\end{aligned} \quad (2)$$

and

$$\begin{aligned}\tilde{a}_{ij}(r) &= \frac{(-1)^j}{n-1} \left((\lambda_2^r T_{i-1}(m_2) T_{j-1}(m_2) + \lambda_3^r T_{i-1}(m_3) T_{j-1}(m_3)) \right. \\ &\quad \left. + 2 \sum_{\substack{k=1 \\ k \neq 2,3}}^n \lambda_k^r T_{i-1}(m_k) T_{j-1}(m_k) \right); \quad i = 1, \dots, n; \quad j = 2, \dots, n-1\end{aligned} \quad (3)$$

where $m_k = \frac{\lambda_k - a}{2b}$ for λ_k is the k -th eigenvalue of $\tilde{A}_n$ see [14, p. 3] and $T_j(\cdot)$ is the j -th degree Chebyshev polynomial of the first kind [11, p. 14]. Öteleş et al. computed integer powers of certain complex tridiagonal matrix

$$\tilde{B}_n = \begin{cases} \tilde{b}_{11} = \tilde{b}_{nn} = a + b \\ \tilde{b}_{12} = \tilde{b}_{21} = \tilde{b}_{n-1,n} = \tilde{b}_{n,n-1} = b & [9, p.67] \\ \text{tridiag}(b, a, b), \text{ other.} \end{cases}$$

as

$$\tilde{b}_{ij}^r = \sum_{k=1}^n f_k \lambda_k^r T_{\frac{2i-1}{2}} \left(\frac{\lambda_k - a}{2b} \right) T_{\frac{2j-1}{2}} \left(\frac{\lambda_k - a}{2b} \right); \quad i, j = 1, \dots, n \quad (4)$$

where $T_j(\cdot)$ is the j -th degree Chebyshev polynomial of the first kind [11, p. 14], λ_k is the k -th eigenvalue of the matrix $\tilde{B}_n$ and

$$f_k = \begin{cases} \frac{2}{n}, & \text{if } k = 1, \dots, n-1 \\ \frac{1}{n}, & \text{if } k = n. \end{cases}$$

Let

$$A_n := \begin{pmatrix} & & & 2b & a \\ & & & -b & a \\ & & \ddots & a & -b \\ & -b & \ddots & \ddots & \\ b & a & -b & & \\ a & 2b & & & \end{pmatrix} \quad (5)$$

and

$$B_n := \begin{pmatrix} & & & b & a+b \\ & & & b & a \\ & & & a & b \\ & \ddots & \ddots & \ddots & \\ b & a & b & & \\ a+b & b & & & \end{pmatrix} \quad (6)$$

be the anti-tridiagonal matrices, where $b \neq 0$ and $a, b \in \mathbb{C}$. In this paper, we obtain the integer powers of the $n \times n$ complex anti-tridiagonal matrices in (5) and (6). We also get the complex factorizations of Fibonacci polynomials, Fibonacci and Pell numbers using the eigenvalues of the matrices A_n and B_n .

2 General Expression of A_n^r

In this section, we obtain a general expression for the entries of the r -th powers of an $n \times n$ complex anti-tridiagonal matrix A_n and B_n in (5) and (6) where if n is even, $r \in \mathbb{N}$ or n is odd, $r \in \mathbb{Z}$.

Lemma 1 *Let $a, 0 \neq b \in \mathbb{C}$, $n \in \mathbb{N}$ and*

$$\tilde{A}_n = \begin{pmatrix} a & 2b & & & \\ b & a & b & & \\ & -b & a & -b & \\ & & -b & \ddots & \ddots \\ & & & \ddots & a & -b \\ & & & & -b & a & b \\ & & & & & 2b & a \end{pmatrix}, \quad (7)$$

$$\tilde{B}_n = \begin{pmatrix} a+b & b & & & \\ b & a & b & & \\ & b & a & b & \\ & & b & \ddots & \ddots \\ & & & \ddots & a & b \\ & & & & b & a & b \\ & & & & & b & a+b \end{pmatrix} \quad (8)$$

and

$$J_n = \begin{pmatrix} & & & & 1 \\ & & & 1 & \\ & & \ddots & & \\ & 1 & & & \\ 1 & & & & \end{pmatrix}. \quad (9)$$

Then

$$A_n = J_n \tilde{A}_n = \tilde{A}_n J_n$$

and

$$B_n = J_n \tilde{B}_n = \tilde{B}_n J_n$$

Proof. See [7]. ■

Lemma 2 Let A_n be an $n \times n$ complex anti-tridiagonal matrix given by (5). Then

$$A_n^r = \begin{cases} J_n \tilde{A}_n^r, & r \text{ is odd} \\ \tilde{A}_n^r, & r \text{ is even.} \end{cases} \quad (10)$$

Proof. We will show by induction on r . The case $r = 1$ is clear. Suppose that the equality (10) is true for $r > 1$. Now let us show the equality (10) is true for $r + 1$. By the induction hypothesis we have

$$A_n^{r+1} = \begin{cases} \tilde{A}_n J_n \tilde{A}_n^r, & r + 1 \text{ is odd} \\ \tilde{A}_n J_n \tilde{A}_n^r J_n, & r + 1 \text{ is even.} \end{cases}$$

Since $A_n = \tilde{A}_n J_n$, we obtain

$$A_n^{r+1} = \begin{cases} J_n \tilde{A}_n^{r+1}, & r + 1 \text{ is odd} \\ \tilde{A}_n^{r+1}, & r + 1 \text{ is even.} \end{cases}$$

■

The same proof can be done easily for the matrix B_n .

Theorem 3 Let A_n be an $n \times n$ complex anti-tridiagonal matrix given by (5). If n is odd, then the r -th power of A_n is

$$\begin{aligned} a_{n-i+1,j}^r &= \frac{1}{2n-2} (\lambda_2^r T_{n-i}(m_2) T_{j-1}(m_2) + \lambda_3^r T_{n-i}(m_3) T_{j-1}(m_3) \\ &\quad + 2 \sum_{\substack{k=1 \\ k \neq 2,3}}^n \lambda_k^r T_{n-i}(m_k) T_{j-1}(m_k)); \quad i = 1, \dots, n; \quad j = 1, n \end{aligned} \quad (11)$$

and

$$\begin{aligned} a_{n-i+1,j}^r &= \frac{(-1)^j}{n-1} \left(\lambda_2^r T_{n-i}(m_2) T_{j-1}(m_2) + \lambda_3^r T_{n-i}(m_3) T_{j-1}(m_3) \right. \\ &\quad \left. + 2 \sum_{\substack{k=1 \\ k \neq 2,3}}^n \lambda_k^r T_{n-i}(m_k) T_{j-1}(m_k) \right) \end{aligned} \quad (12)$$

for $i = 1, \dots, n; \quad j = 2, \dots, n-1$ and if n is even, then the r -th power of A_n is

$$\begin{aligned} a_{i,j}^r &= \frac{1}{2n-2} (\lambda_2^r T_{i-1}(m_2) T_{j-1}(m_2) + \lambda_3^r T_{i-1}(m_3) T_{j-1}(m_3) \\ &\quad + 2 \sum_{\substack{k=1 \\ k \neq 2,3}}^n \lambda_k^r T_{i-1}(m_k) T_{j-1}(m_k)); \quad i = 1, \dots, n; \quad j = 1, n \end{aligned} \quad (13)$$

and

$$a_{i,j}^r = \frac{(-1)^j}{n-1} \left(\lambda_2^r T_{i-1}(m_2) T_{j-1}(m_2) + \lambda_3^r T_{i-1}(m_3) T_{j-1}(m_3) + 2 \sum_{\substack{k=1 \\ k \neq 2,3}}^n \lambda_k^r T_{i-1}(m_k) T_{j-1}(m_k) \right) \quad (14)$$

for $i = 1, \dots, n$; $j = 2, \dots, n-1$.

Proof. Let $A_n^r = (a_{i,j}^r)$ and $U = \tilde{A}_n^r$. We obtained the eigenvalues of $\tilde{A}_n$ as

$$\lambda_k = a + 2b \cos \left(\frac{(k-1)\pi}{n-1} \right), \quad k = 1, \dots, n \quad (\text{see [14, p.3]})$$

and the entries of the matrix U as

$$u_{ij}(r) = \frac{1}{2n-2} (\lambda_2^r T_{i-1}(m_2) T_{j-1}(m_2) + \lambda_3^r T_{i-1}(m_3) T_{j-1}(m_3) + 2 \sum_{\substack{k=1 \\ k \neq 2,3}}^n \lambda_k^r T_{i-1}(m_k) T_{j-1}(m_k)); \quad i = 1, \dots, n; \quad j = 1, n \quad (15)$$

and for $i = 1, \dots, n$ and $j = 2, \dots, n-1$

$$u_{ij}(r) = \frac{(-1)^j}{n-1} \left(\lambda_2^r T_{i-1}(m_2) T_{j-1}(m_2) + \lambda_3^r T_{i-1}(m_3) T_{j-1}(m_3) + 2 \sum_{\substack{k=1 \\ k \neq 2,3}}^n \lambda_k^r T_{i-1}(m_k) T_{j-1}(m_k) \right) \quad (16)$$

where $m_k = \frac{\lambda_k - a}{2b}$ (see [14, p.]) and $T_s(\cdot)$ is the s -th degree Chebyshev polynomial of the first kind [13, p. 14]. If r is even, then the equalities (15) and (16) are valid by the equality (10). Let r be odd number. If we multiply the equalities (15) and (16) by J_n from left side, then we have

$$(J_n \tilde{A}_n^r)_{i,k} = \sum_{s=1}^n (J_n)_{i,s} u_{s,k}(r) = u_{n-i+1,k}(r); \quad k = 1, \dots, n.$$

Hence we obtain

$$\begin{aligned} a_{n-i+1,j}^r &= u_{n-i+1,j}(r) \\ &= \frac{1}{2n-2} (\lambda_2^r T_{n-i}(m_2) T_{j-1}(m_2) + \lambda_3^r T_{n-i}(m_3) T_{j-1}(m_3) + 2 \sum_{\substack{k=1 \\ k \neq 2,3}}^n \lambda_k^r T_{n-i}(m_k) T_{j-1}(m_k)); \quad i = 1, \dots, n; \quad j = 1, n \end{aligned}$$

and

$$\begin{aligned}
a_{n-i+1,j}^r &= u_{n-i+1,j}(r) \\
&= \frac{(-1)^j}{n-1} \left(\lambda_2^r T_{n-i}(m_2) T_{j-1}(m_2) + \lambda_3^r T_{n-i}(m_3) T_{j-1}(m_3) \right. \\
&\quad \left. + 2 \sum_{\substack{k=1 \\ k \neq 2,3}}^n \lambda_k^r T_{n-i}(m_k) T_{j-1}(m_k) \right); i = 1, \dots, n; j = 2, \dots, n-1.
\end{aligned}$$

■

Theorem 4 Let B_n be an $n \times n$ complex anti-tridiagonal matrix given by (6), $\tilde{V} = \tilde{B}_n^r$ and $V = B_n^r$. Then the entries of V are

$$v_{ij}(r) = \begin{cases} \sum_{k=1}^n f_k \lambda_k^r T_{\frac{2i-1}{2}} \left(\frac{\lambda_k - a}{2b} \right) T_{\frac{2j-1}{2}} \left(\frac{\lambda_k - a}{2b} \right), & \text{if } r \text{ is even} \\ \sum_{k=1}^n f_k \lambda_k^r T_{\frac{2(n-i)+1}{2}} \left(\frac{\lambda_k - a}{2b} \right) T_{\frac{2j-1}{2}} \left(\frac{\lambda_k - a}{2b} \right), & \text{if } r \text{ is odd} \end{cases} \quad (17)$$

where $i, j = 1, \dots, n$ and

$$\lambda_k = a + 2b \cos \left(\frac{(k-1)\pi}{n} \right), \quad k = 1, \dots, n$$

are the eigenvalues of the matrix $\tilde{B}_n$.

Proof. Since $B_n^r = \tilde{B}_n^r$ from (10) for r is even, the equality (17) is valid. Let r be odd number. If we multiply the equality (4) by J_n from the left side, then we have

$$\begin{aligned}
v_{i,j} &= (J_n \tilde{B}_n^r)_{i,j} = \sum_{s=1}^n (J_n)_{i,s} \tilde{v}_{s,k}(r) \\
&= \tilde{v}_{n-i+1,j}(r) \\
&= \sum_{k=1}^n f_k \lambda_k^r T_{\frac{2(n-i)+1}{2}} \left(\frac{\lambda_k - a}{2b} \right) T_{\frac{2j-1}{2}} \left(\frac{\lambda_k - a}{2b} \right).
\end{aligned}$$

■

Corollary 5 Let the matrix A_n be as in (5). Then the eigenvalues of A_n are

$$\lambda_k = (-1)^k \left(a + 2b \cos \left(\frac{(k-1)\pi}{n-1} \right) \right), \quad k = 1, \dots, n. \quad (18)$$

Proof. See [3, p.574] and [14, p. 5]. ■

Corollary 6 *Let the matrix B_n be as in (6). Then the eigenvalues of B_n are*

$$\lambda_k = (-1)^{k-1} \left(a + 2b \cos \left(\frac{(k-1)\pi}{n} \right) \right) \quad (19)$$

where $k = 1, \dots, n$.

Proof. See [5, p.574] and [14, p.]. ■

3 Numerical examples

Considering the Eqs. (11-14), we can find the arbitrary integer powers of the $n \times n$ complex anti-tridiagonal matrix A_n in (5), where n is positive odd integer.

Example 7 *Let $n = 3, r = 3, a = 1$ and $b = 3$. Since*

$$\tilde{J} = \text{diag}(\lambda_1, \lambda_2, \lambda_3) = \text{diag}(a, a + 2b, a - 2b) = \text{diag}(1, 7, -5)$$

and

$$\tilde{A}_3^3 = (u_{ij}(3)) = \begin{pmatrix} 55 & 234 & 54 \\ 117 & 109 & 117 \\ 54 & 234 & 55 \end{pmatrix},$$

we obtain

$$A_3^3 = J_3 \tilde{A}_3^3 = \begin{pmatrix} 54 & 234 & 55 \\ 117 & 109 & 117 \\ 55 & 234 & 54 \end{pmatrix}.$$

Example 8 *If $n = 5, r = 4, a = 1$ and $b = 3$, then*

$$\begin{aligned} \tilde{J} &= \text{diag}(\lambda_1, \lambda_2, \lambda_3, \lambda_4, \lambda_5) \\ &= \text{diag}(1, 7, -5, 1 + 3\sqrt{2}, 1 - 3\sqrt{2}) \end{aligned}$$

and since

$$\tilde{A}_5^4 = (u_{ij}(4)) = \begin{pmatrix} 595 & 672 & -756 & 216 & 162 \\ 336 & 973 & -444 & 540 & 108 \\ -378 & -444 & 757 & -444 & -378 \\ 108 & 540 & -444 & 973 & 336 \\ 162 & 216 & -756 & 672 & 595 \end{pmatrix},$$

we have

$$A_5^4 = J_5 \tilde{A}_5^4 = \begin{pmatrix} 162 & 216 & -756 & 672 & 595 \\ 108 & 540 & -444 & 973 & 336 \\ -378 & -444 & 757 & -444 & -378 \\ 336 & 973 & -444 & 540 & 108 \\ 595 & 672 & -756 & 216 & 162 \end{pmatrix}.$$

4 Complex Factorizations

The well-known the Fibonacci polynomials $F(x) = \{F_n(x)\}_{n=1}^{\infty}$ are defined by $F_n(x) = xF_{n-1}(x) + F_{n-2}(x)$ with initial conditions $F_0(x) = 0$ and $F_1(x) = 1$. For $x = 1$ and $x = 2$, then we obtain the Fibonacci and Pell numbers as

$$F_n(1) = \{0, 1, 1, 2, 3, 5, 8, \dots\}$$

and

$$F_n(2) = \{0, 1, 2, 5, 12, 29, \dots\},$$

respectively.

Theorem 9 *Let A_n be an $n \times n$ complex anti-tridiagonal matrix given by (5). If $a := x$ and $b := \mathbf{i}$, then*

$$\det(A_n) = \begin{cases} (x^2 + 4)F_{n-1}(x), & n \equiv 0 \text{ or } 1 \pmod{4} \\ -(x^2 + 4)F_{n-1}(x), & n \equiv 2 \text{ or } 3 \pmod{4} \end{cases}$$

where $\mathbf{i} = \sqrt{-1}$.

Proof. Since $\tilde{A}_n = J_n A_n$,

$$\det(\tilde{A}_n) = x^2 D_{n-2} + 4x D_{n-3} + 4D_{n-4}$$

where $D_n = \det(\text{tridiag}_n(-\mathbf{i}, x, -\mathbf{i}))$ and

$$\det(\text{tridiag}_n(-\mathbf{i}, x, -\mathbf{i})) = F_{n+1}(x) \text{ (see [14, p.]},$$

we arrive at

$$\begin{aligned} \det(\tilde{A}_n) &= x^2 F_{n-1}(x) + 4x F_{n-2}(x) + 4F_{n-3}(x) \\ &= x^2 (x F_{n-2}(x) + F_{n-3}(x)) + 4x F_{n-2}(x) + 4F_{n-3}(x) \\ &= (x^2 + 4)(x F_{n-2}(x) + F_{n-3}(x)) = (x^2 + 4)F_{n-1}(x). \end{aligned}$$

Since

$$\det(J_n) = \begin{cases} 1, & n \equiv 0 \text{ or } 1 \pmod{4} \\ -1, & n \equiv 2 \text{ or } 3 \pmod{4}, \end{cases}$$

the proof of theorem is completed. ■

Corollary 10 *Let A_n be an $n \times n$ complex anti-tridiagonal matrix given by (5). If $a := x$ and $b := \mathbf{i}$, then the complex factorization of generalized Fibonacci-Pell numbers is the following form:*

$$F_{n-1}(x) = \frac{\alpha}{x^2 + 4} \prod_{k=1}^n \left(x + 2\mathbf{i} \cos \left(\frac{(k-1)\pi}{n-1} \right) \right)$$

where

$$\alpha = \begin{cases} (-1)^n, & \text{if } n \equiv 0 \text{ or } 1 \pmod{4} \\ (-1)^{n-1}, & \text{if } n \equiv 2 \text{ or } 3 \pmod{4}. \end{cases}$$

Proof. Since the eigenvalues of the matrix A_n from (18)

$$\lambda_j = (-1)^j \left(x + 2\mathbf{i} \cos \left(\frac{(j-1)\pi}{n-1} \right) \right), \quad j = 1, \dots, n,$$

the determinant of the matrix A_n can be obtained as

$$\det(A_n) = (-1)^n \prod_{k=1}^n \left(x + 2\mathbf{i} \cos \left(\frac{(k-1)\pi}{n-1} \right) \right). \quad (20)$$

By considering (20) and Theorem 9, the complex factorization of generalized Fibonacci-Pell numbers is obtained. ■

Theorem 11 *Let B_n be an $n \times n$ complex anti-tridiagonal matrix given by (6). If $a := 1$ and $b := \mathbf{i}$, then*

$$\det(B_n) = \begin{cases} (1 + 2\mathbf{i})F_n, & n \equiv 0 \text{ or } 1 \pmod{4} \\ -(1 + 2\mathbf{i})F_n, & n \equiv 2 \text{ or } 3 \pmod{4} \end{cases} \quad (21)$$

and if $a := 2$ and $b := \mathbf{i}$, then

$$\det(B_n) = \begin{cases} (2 + 2\mathbf{i})P_n, & n \equiv 0 \text{ or } 1 \pmod{4} \\ -(2 + 2\mathbf{i})P_n, & n \equiv 2 \text{ or } 3 \pmod{4} \end{cases} \quad (22)$$

where $\mathbf{i} = \sqrt{-1}$ and F_n and P_n denote the n th Fibonacci and Pell numbers, respectively.

Proof. Applying Laplace expansion according to the first two and last two rows of the determinant of $\tilde{B}_n$, we have

$$\begin{aligned} \det(\tilde{B}_n) &= (a+b)^2 \det(\text{tridiag}_{n-2}(b, a, b)) \\ &\quad - 2b^2(a+b) \det(\text{tridiag}_{n-3}(b, a, b)) \\ &\quad + b^4 \det(\text{tridiag}_{n-4}(b, a, b)) \quad (\text{see [9, p.]}). \end{aligned} \quad (23)$$

If we take $a := 1$ and $b := \mathbf{i}$ in (23), then we get

$$\begin{aligned} \det(\tilde{B}_n) &= (1 + \mathbf{i})^2 \det(\text{tridiag}_{n-2}(\mathbf{i}, 1, \mathbf{i})) \\ &\quad + 2(1 + \mathbf{i}) \det(\text{tridiag}_{n-3}(\mathbf{i}, 1, \mathbf{i})) \\ &\quad + \det(\text{tridiag}_{n-4}(\mathbf{i}, 1, \mathbf{i})) \quad (\text{see [9, p.]}) \\ &= (1 + \mathbf{i})^2 F_{n-1} + 2(1 + \mathbf{i}) F_{n-2} + F_{n-3} \\ &= (1 + 2\mathbf{i}) F_n. \end{aligned}$$

Since

$$B_n = J_n \tilde{B}_n,$$

we obtain (21).

Similar to the above, we can easily obtain Pell numbers. ■

Corollary 12 Let B_n be an $n \times n$ complex anti-tridiagonal matrix given by (6). If $a := 1$ and $b := \mathbf{i}$, then the complex factorizations of Fibonacci numbers are

$$F_n = (-1)^n \begin{cases} \prod_{k=2}^{n-1} \left(1 + 2\mathbf{i} \cos \left(\frac{(k-1)\pi}{n} \right) \right), & n \text{ is even} \\ - \prod_{k=2}^n \left(1 + 2\mathbf{i} \cos \left(\frac{(k-1)\pi}{n} \right) \right), & n \text{ is odd} \end{cases}$$

and if $a := 2$ and $b := \mathbf{i}$, then the complex factorizations of Pell numbers are

$$P_n = (-1)^n \begin{cases} \prod_{k=2}^n \left(2 + 2\mathbf{i} \cos \left(\frac{(k-1)\pi}{n} \right) \right), & n \text{ is even} \\ - \prod_{k=2}^n \left(2 + 2\mathbf{i} \cos \left(\frac{(k-1)\pi}{n} \right) \right), & n \text{ is odd.} \end{cases}$$

Proof. Let $a := 1$ and $b := \mathbf{i}$. Since the eigenvalues of the matrix B_n are

$$\lambda_k = (-1)^{n-1} \left(1 + 2\mathbf{i} \cos \left(\frac{(k-1)\pi}{n} \right) \right)$$

for $k = 1, \dots, n$ and the determinant of the matrix B_n is equal to multiplication of its eigenvalues, we get

$$\begin{aligned} F_n &= \frac{1}{1 + 2\mathbf{i}} \begin{cases} \det(B_n), & n \equiv 0 \text{ or } 1 \pmod{4} \\ -\det(B_n), & n \equiv 2 \text{ or } 3 \pmod{4} \end{cases} \\ &= \frac{1}{1 + 2\mathbf{i}} \begin{cases} (-1)^n \prod_{k=1}^n \left(1 + 2\mathbf{i} \cos \left(\frac{(k-1)\pi}{n} \right) \right), & n \text{ is even} \\ (-1)^{n+1} \prod_{k=1}^n \left(1 + 2\mathbf{i} \cos \left(\frac{(k-1)\pi}{n} \right) \right), & n \text{ is odd} \end{cases} \\ &= (-1)^n \begin{cases} \prod_{k=2}^n \left(1 + 2\mathbf{i} \cos \left(\frac{(k-1)\pi}{n} \right) \right), & n \text{ is even} \\ - \prod_{k=2}^n \left(1 + 2\mathbf{i} \cos \left(\frac{(k-1)\pi}{n} \right) \right), & n \text{ is odd.} \end{cases} \end{aligned}$$

We can easily obtain Pell numbers similarly for $a := 2$ and $b := \mathbf{i}$.

Thus, the proof is completed. ■

Acknowledgement. The authors are partially supported by TUBITAK and the Office of Selçuk University Research Project (BAP).

References

- [1] J. Rimas, Integer powers of real odd order skew-persymmetric anti-tridiagonal matrices with constant anti-diagonals ($\text{antitridiag}_n(a, c, -a)$, $a \in \mathbb{R} - \{0\}$, $c \in \mathbb{R}$), Applied Mathematics and Computation, 219, (2013) 7075-7088.

- [2] J. Rimas, Integer powers of real even order anti-tridiagonal Hankel matrices of the form $\text{antitridiag}_n(a, c, -a)$, Applied Mathematics and Computation, 225, (2013) 204-215.
- [3] J. Rimas, On computing of arbitrary positive integer powers for one type of symmetric anti-tridiagonal matrices of odd order, Applied Mathematics and Computation, 203, (2008) 573-581.
- [4] J. Rimas, On computing of arbitrary positive integer powers for one type of tridiagonal matrices, Applied Mathematics and Computation, 161, (2005) 1037-1040.
- [5] J. Rimas, On computing of arbitrary positive integer powers for one type of tridiagonal matrices with elements $1, 0, 0, \dots, 0, 1$ in principal and $1, 1, \dots, 1$ in neighboring diagonals -I, Applied Mathematics and Computation, 186, (2007) 1254-1257.
- [6] Q. Yin, On computing of arbitrary positive powers for anti-tridiagonal matrices of even order, Applied Mathematics and Computation, 203, (2008) 252-257.
- [7] H. Wang, Powers of complex persymmetric antitridiagonal matrices with constant antidiagonals, Computational Mathematics, 2014 (2014) 1-10.
- [8] A. Öteleş & M. Akbulak, Positive integer powers of certain complex tridiagonal matrices, Applied Mathematics and Computation, 219, (2013) 10448-10455.
- [9] A. Öteleş & M. Akbulak, Positive integer powers of certain complex tridiagonal matrices, Mathematical Science Letters, 2-1 (2013) 63-72.
- [10] A. Öteleş & M. Akbulak, Positive integer powers of one type of complex tridiagonal matrix, Bulletin of the Malaysian Mathematical Science Society, 2, 37(3) 2014, 1-xx.
- [11] Jesus Gutiérrez-Gutiérrez, Powers of complex persymmetric or skew-persymmetric anti-tridiagonal matrices with constant anti-diagonals, Applied Mathematics and Computation, 217, (2011) 6125-6132.
- [12] Honglin Wu, On computing of arbitrary positive powers for one type of anti-tridiagonal matrices of even order, Applied Mathematics and Computation, 217, (2010) 2750-2756.
- [13] J. C. Mason & D. C. Handscomb, Chebyshev Polynomials, CRC Press, Washington, 2003.
- [14] D. Bozkurt & Ş. B. Bozkurt, Integer powers of certain complex tridiagonal matrices and some complex factorizations, <http://arxiv.org/pdf/1403.6696v1.pdf>.